


How to dissolve and drink Magcorol Powder 50 g

How to Dissolve

①



Fill a large glass with 150mL water.

②



Add all the MAGCOROL Powder while stirring with a spoon, and mix well.

Once you put MAGCOROL Powder in water, do not leave it unattended; dissolve it immediately.

How to Drink

- Take your time and take each sip slowly.
- Elderly people and people who had their stomachs removed should take the whole dose by taking time especially slowly.
- Please drink as much water as possible before and after taking the dose.
- Do not take this dose alone.
- Add 2-3 pieces of ice to make it colder, making it easier to drink.
- * You will have your first bowel movement approximately 2-10 hours after you start drinking.
(There are individual differences in defecation.)
- * After taking the dose, you will have watery stools for several times.
- * If your stool volume is low or your stools are not watery, please inform us before the undergoing the examination.

[Caution] Please contact us if you have experienced:

- No bowel movements either on the day before or before taking MAGCOROL Powder
(in this case, do not take the dose and contact us);
- No bowel movements after taking a full dose of MAGCOROL Powder; or
- Any symptoms occurring while or after taking MAGCOROL Powder including:
 - ✓ Feel pain in the belly
 - ✓ Feel like throwing up
 - ✓ Threw up
 - ✓ Had bloody stool
 - ✓ Feel ill
 - ✓ Feel tired
 - ✓ Feel drowsy
 - ✓ Have skin redness
 - ✓ Got hives